



## PATENT ABSTRACTS OF JAPAN

(11) Publication number: 09048770 A

(43) Date of publication of application: 18.02.1997

(51) Int. Cl. C07D263/44

A61K 31/41, A61K 31/42, A61K 31/42, A61K 31/425, A61K 31/425,  
 A61K 31/425, A61K 31/425, A61K 31/425, A61K 31/44, A61K 31/47,  
 A61K 31/495, C07D271/06, C07D277/34, C07D413/12, C07D413/12,  
 C07D413/12, C07D413/12, C07D413/14, C07D413/14, C07D417/12,  
 C07D417/12, C07D417/12, C07D417/12, C07D417/12, C07D417/12,  
 C07D417/12, C07D417/12, C07D417/12, C07D417/14, C07D417/14,  
 C07D417/14, C07D417/14, C07D417/14, C07D417/14, C07D417/14,  
 C07D493/04, C07D493/04

(21) Application number: 08136609  
 (22) Date of filing: 30.05.1996  
 (30) Priority: 02.06.1995 JP 07137002

(71) Applicant: SANKYO CO LTD  
 (72) Inventor: YANAGISAWA HIROAKI  
 FUJITA TAKESHI  
 FUJIMOTO KOICHI  
 WADA KUNIO  
 OGUCHI MINORU  
 YOSHIOKA TAKAO  
 FUJIWARA TOSHIHIKO  
 HORIKOSHI HIROYOSHI

(54) AROMATIC OXYIMINO DERIVATIVE


lowed by deprotecting the resultant compound.

COPYRIGHT: (C)1997,JPO


(57) Abstract:

PROBLEM TO BE SOLVED: To obtain an aromatic oxyimino derivative having remedying effect on insulin resistance, hyperlipemia, hyperglycemia, obesity, etc., excellent in safety, thus useful for preventing and treating these diseases.


SOLUTION: This new compound (salt) is expressed by formula I [ $R^1$  is H, a 1-6C alkyl, 1-4C alkoxy, 1-4C alkylthio, etc.;  $R^2$  is a 1-6C alkylene; A is methylene, O, S, etc.; B is (1- $3R^1$  (except H)-substituted) 5- or 6-membered heterocyclic divalent group, etc.; X is (2,4,5-trihydro-3H-2,4-dioxo-thiazol-5-yl) methyl, etc.; Y is S, O, etc.], e.g. 5-(4-{2-[(6,7-dichloroindan-1-yl) iminoxy]ethoxy}benzyl)-2,4-thiazolidindione. The compound of formula I is obtained by condensation reaction between a compound of formula II (U is OH, halogen, etc.) and a compound of formula III ( $X'$  is a group resulted by protecting the NH group at X) fol-



I



II



III